

Durham Research Online

Deposited in DRO:

21 August 2015

Version of attached file:

Accepted Version

Peer-review status of attached file:

Not peer-reviewed

Citation for published item:

Paulusma, D. (2016) 'Open problems on graph coloring for special graph classes.', in Graph-theoretic concepts in computer science : 41st international workshop, WG 2015, Garching, Germany, June 17-19, 2015 ; revised papers. Berlin: Springer, pp. 16-30. Lecture notes in computer science. (9224).

Further information on publisher's website:

http://dx.doi.org/10.1007/978-3-662-53174-7_2

Publisher's copyright statement:

The final publication is available at Springer via http://dx.doi.org/10.1007/978-3-662-53174-7_2

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Open Problems on Graph Coloring for Special Graph Classes

Daniël Paulusma^{*}

School of Engineering and Computing Sciences, Durham University
Science Laboratories, South Road, Durham DH1 3LE, UK
`daniel.paulusma@durham.ac.uk`

Abstract. For a given graph G and integer k , the COLORING problem is that of testing whether G has a k -coloring, that is, whether there exists a vertex mapping $c : V \rightarrow \{1, 2, \dots\}$ such that $c(u) \neq c(v)$ for every edge $uv \in E$. We survey known results on the computational complexity of COLORING for graph classes that are hereditary or for which some graph parameter is bounded. We also consider coloring variants, such as precoloring extensions and list colorings and give some open problems in the area of on-line coloring.

1 Introduction

Graph coloring is a central topic in Computer Science due to a high number of theoretical and practical applications. Within both structural and algorithmic graph theory, many graph coloring variants and generalizations have been studied. Besides the well-known text-book of Toft and Jensen [65], several survey papers appeared over the years. For instance, the survey of Tuza [70] considered the graph coloring problem and variants of it, in which local restrictions are imposed on the coloring (e.g. precoloring extensions and list colorings) whereas the survey of Randerath and Schiermeyer [61] considered structural and complexity aspects of graph colorings for hereditary graph classes.

As graph coloring and many of its variants are computationally hard on general graphs, it is natural to restrict the input graph to some special graph class. This topic has been extensively studied in the literature. A recent survey of Golovach, Johnson, Paulusma and Song [27] updated several parts of the two aforementioned survey papers [61, 70] and the survey of Chudnovsky [13] by primarily focussing on computational complexity aspects of graph coloring for graph classes characterized by one or two forbidden induced subgraphs. As noted by Golovach et al. [27], the task to collect complexity results for graph coloring restricted to other graph classes might be beyond the scope of a single paper. Our aim is therefore to discuss a number of such results and open problems not mentioned in [27]. In particular we will identify a number of gaps in existing complexity results.

^{*} The author was supported by EPSRC (EP/K025090/1).

The survey is organized as follows. In Section 2 we state some terminology. Then, in Section 3, we consider the (classical) complexity of graph coloring, pre-coloring extension and list colorings for a number of graph classes characterized by more than two forbidden induced subgraphs and also for some graph classes for which some graph parameter is bounded. We discuss parameterized coloring problems in Section 4 and on-line coloring problems in Section 5. We briefly consider graph homomorphisms in Section 6.




Figure 1. Relationships between COLORING and its variants as shown in [27]. An arrow from one problem to another indicates that the latter is a special case of the former; k and ℓ are any two integers for which $\ell \geq k$.

2 Preliminaries

A *coloring* of a graph $G = (V, E)$ is a vertex mapping $c : V \rightarrow \{1, 2, \dots\}$ with the additional condition that $c(u) \neq c(v)$ whenever $uv \in E$. We call $c(u)$ the *color* of u . If $1 \leq c(u) \leq k$ for all $u \in V$ then c is also called a *k-coloring* of G . We say that G is *k-colorable* if a k -coloring of G exists. The *chromatic number* $\chi(G)$ of G is the smallest integer k for which G is k -colorable. The COLORING problem is that of deciding whether a graph G is k -colorable for some given integer k . If k is fixed (that is, not part of the input) we obtain the k -COLORING problem.

A *k-precoloring* of a graph $G = (V, E)$ is a mapping $c_W : W \rightarrow \{1, 2, \dots, k\}$ for some subset $W \subseteq V$. We say that a k -coloring c of G is an extension of a k -precoloring c_W of G if $c(v) = c_W(v)$ for each $v \in W$. For a given graph G , a positive integer k and a k -precoloring c_W of G , the PRECOLORING EXTENSION problem asks whether c_W can be extended to a k -coloring of G . If k is fixed we denote this problem as the k -PRECOLORING EXTENSION problem.

A *list assignment* of a graph $G = (V, E)$ is a function L with domain V such that for each vertex $u \in V$, $L(u)$ is a subset of $\{1, 2, \dots\}$. This set is called the *list of admissible colors* for u . If $L(u) \subseteq \{1, \dots, k\}$ for each $u \in V$ then L is also called a *k-list assignment*. The *size* of a list assignment L is the maximum list size $|L(u)|$ over all vertices $u \in V$. A coloring c *respects* L if $c(u) \in L(u)$ for

all $u \in V$. This leads to the following three problems. Given a graph G with a list assignment L , the LIST COLORING problem is that of testing whether G has a coloring that respects L . If inputs are restricted to pairs (G, L) where L has size at most ℓ then we obtain the ℓ -LIST COLORING problem, and if each L is a k -list assignment, we obtain the LIST k -COLORING problem. See Figure 1 for a display of the relationships between the seven problems defined above.

Let G be a graph and $\{H_1, \dots, H_p\}$ be a set of graphs. Then G is said to be (H_1, \dots, H_p) -free if G has no *induced* subgraph isomorphic to a graph in $\{H_1, \dots, H_p\}$. If $p = 1$, we may write that G is H_1 -free. The disjoint union $(V(G) \cup V(H), E(G) \cup E(H))$ of two vertex-disjoint graphs G and H is denoted by $G + H$ and the disjoint union of r copies of a graph G is denoted by rG . The complement \overline{G} of a graph G has vertex set $V(\overline{G}) = V(G)$ and an edge between two distinct vertices u and v if and only if u and v are not adjacent in G . We denote the path and cycle on n vertices by P_n and C_n , respectively.

Let G be a connected graph. The *distance* $d(u, v)$ between two vertices u and v in G is the number of edges in a shortest path from u to v in G . The *diameter* of G is the maximum of $\max_v d(u, v)$ over all vertices u in G . The *radius* of G is the minimum of $\max_v d(u, v)$ over all vertices u in G .

Let p be a graph parameter and let \mathcal{G} be a graph class. We say that \mathcal{G} has *bounded* p if there exists a constant c such that $p(G) \leq c$ for all $G \in \mathcal{G}$.

3 Classical Complexity

The problems k -COLORING, k -PRECOLORING EXTENSION, LIST k -COLORING and k -LIST COLORING are polynomial-time solvable for general graphs if $k \leq 2$ and NP-complete if $k \geq 3$ [52, 67]. As mentioned, we refer to the recent survey [27] for an overview of known results for these problems when restricted to H -free graphs and (H_1, H_2) -free graphs. In this section we consider a number of other graph classes.

Cycle-free Graphs. A *hole* is a cycle of on at least four vertices. An *antihole* is the complement of a hole. A cycle, hole or antihole is *even* if it contains an even number of vertices; otherwise it is *odd*. An (anti)hole is *long* if it has at least five vertices. A graph is *odd-hole-free* or *odd-antihole-free* if it contains no induced odd holes or no induced odd antiholes, respectively. In a similar way we define (even-)hole-free, (even-)antihole-free, long-hole-free, long-antihole-free, odd-cycle-free and (odd-)anticycle-free graphs.

Grötschel, Lovász, and Schrijver, [30] proved that COLORING is polynomial-time solvable for perfect graphs, or equivalently (due to the Strong Perfect Graph Theorem [14]) for graphs that are odd-hole-free and odd-antihole-free. Note that hole-free graphs and antihole-free graphs are perfect. Hence COLORING is also polynomial-time solvable for hole-free graphs and antihole-free graphs, and thus for cycle-free graphs (forests) and anticycle-free graphs (coforests).

Král', Kratochvíl, Tuza and Woeginger [49] proved that COLORING is NP-complete for $(2P_2, C_5)$ -free graphs and also for (C_3, C_4, C_5) -free graphs. Consequently, COLORING is NP-complete for long-hole-free graphs and long-antihole-

free graphs, and thus for odd-hole-free graphs and odd-antihole free graphs. In contrast, COLORING is polynomial-time solvable for odd-cycle-free graphs (bipartite graphs) and odd-anticycle-free graphs (cobipartite graphs). If we change the parity of the forbidden cycles from odd to even, then we obtain two long-standing open problems; we refer to the survey of Vušković [68] for more on even-cycle-free graphs (even-hole-free graphs).

Open Problem 1 *Determine the complexity of COLORING for even-cycle-free graphs and even-anticycle-free graphs.*

r	k -PRECOLORING EXT.				LIST k -COLORING			
	$k = 3$	$k = 4$	$k = 5$	$k \geq 6$	$k = 3$	$k = 4$	$k = 5$	$k \geq 6$
$t \leq 6$	P	P	P	P	P	P	P	P
$t = 7$?	?	?	?	?	?	?	?
$t = 8$?	?	?	?	?	NP-c	NP-c	NP-c
$t = 9$?	?	?	?	?	NP-c	NP-c	NP-c
$t \geq 10$?	NP-c	NP-c	NP-c	?	NP-c	NP-c	NP-c

Table 1. The complexity of k -PRECOLORING EXTENSION and LIST k -COLORING on P_t -free bipartite graphs for fixed k and t .

Bipartite and Chordal Bipartite Graphs. A graph is *chordal bipartite* if it is bipartite and every induced cycle has exactly four vertices. Hujter and Tuza [41] proved that PRECOLORING EXTENSION is linear-time solvable on P_5 -free bipartite graphs (which are chordal bipartite) and NP-complete for P_6 -free chordal bipartite graphs. Kratochvíl [50] answered two of their open problems [40] by proving that 3-PRECOLORING EXTENSION is NP-complete for planar bipartite graphs and that 5-PRECOLORING EXTENSION is NP-complete for P_{14} -free bipartite graphs. The latter result was strengthened by Huang et al. [39], who proved that, for all $k \geq 4$, k -PRECOLORING EXTENSION is NP-complete for P_{10} -free chordal bipartite graphs. The same authors [39] also proved that LIST 4-COLORING is NP-complete for P_8 -free chordal bipartite graphs.

Brandstädt, Klemmt and Mahfud [5] proved that the class of (C_3, P_6) -free graphs has bounded clique-width. By combining their result with results of Kobler and Rotics [48] and Oum and Seymour [62] we find that, for all $k \geq 1$, LIST k -COLORING is polynomial-time solvable on (C_3, P_6) -free graphs (see also e.g. [39]). Table 1 summarizes the above results for P_t -free bipartite graphs.

Open Problem 2 *Determine the complexity of the problems k -PRECOLORING EXTENSION and LIST k -COLORING for the missing cases in Table 1.*

Huang et al. [39] posed the following two open problems.

Open Problem 3 *Determine the complexity of the problems LIST 3-COLORING and 3-PRECOLORING EXTENSION for the class of chordal bipartite graphs.*

Planar Graphs and Graphs of Bounded Vertex Degree. To recall two classic results, Garey, Johnson and Stockmeyer [26] proved that 3-COLORING is NP-complete even for planar graphs of maximum degree 4, whereas every planar graph is 4-colorable by the Four Color Theorem [2]. Chlebík and Chlebíková [12] strengthened the aforementioned result of Kratochvíl [50] for planar bipartite graphs by proving that 3-PRECOLORING EXTENSION is NP-complete even for planar bipartite graphs of maximum degree 4. The same authors [12] also showed that LIST 3-COLORING is NP-complete for 3-regular planar bipartite graphs but that PRECOLORING EXTENSION is polynomial-time solvable for arbitrary graphs of maximum degree at most 3.

Demange and de Werra [17] proved that 3-PRECOLORING EXTENSION is NP-complete for subgrids (which are induced subgraphs of grids and hence have maximum degree at most 4) and that LIST 3-COLORING is NP-complete even for subgrids of maximum degree at most 3, whereas Kratochvíl and Tuza [51] showed that LIST COLORING is polynomial-time solvable for graphs of maximum degree 2. Demange and de Werra [17] also proved that LIST 4-COLORING is NP-complete for grids, and they posed the following open problem.

Open Problem 4 *Determine the complexity of PRECOLORING EXTENSION for grids.*

When consider graphs of bounded degree that are not necessarily planar a full complexity classification is known (see [16]) for COLORING, LIST COLORING and PRECOLORING EXTENSION and also for k -LIST COLORING, LIST k -COLORING and k -PRECOLORING EXTENSION. However, no dichotomy is known for k -COLORING restricted to graphs of maximum degree at most d , but some partial results have been obtained. Molloy and Reed [60] classified the complexity for all pairs (k, d) for sufficiently large d , whereas Emden-Weinert et al. [19] showed that k -COLORING is NP-complete for graphs of maximum degree at most $k + \lceil \sqrt{k} \rceil - 1$. By combining the latter result with Brooks' Theorem [8], we find that the smallest open case is the following problem.

Open Problem 5 *Determine the complexity of 5-COLORING for graphs of maximum degree 6.*

Graphs of Bounded Diameter. By using a reduction from 3-COLORING via adding dominating vertices one can easily show that k -COLORING is NP-complete for graphs of diameter d for all pairs (k, d) with $k \geq 3$ and $d \geq 2$ except for two notorious cases, namely $(k, d) \in \{(3, 2), (3, 3)\}$. Mertzios and Spirakis [57] settled the case $(k, d) = (3, 3)$. They proved that, for every $0 \leq \epsilon < 1$, 3-COLORING is NP-complete even for classes of triangle-free graphs $G = (V, E)$ of diameter 3, radius 2 and minimum degree $\delta = \Theta(|V|^\epsilon)$.

We note that, for every $k \geq 1$ and $p \geq 1$, the problems k -COLORING and k -PRECOLORING EXTENSION are polynomially equivalent on the class of graphs of diameter at most p . This can be seen as follows. Firstly, k -COLORING is a special case of k -PRECOLORING EXTENSION. Secondly, if we are given a graph G of diameter at most p with a k -precoloring c_W for some $W \subseteq V(G)$, then we

identify any two vertices of W that are colored alike. Afterwards all precolored vertices have a distinct color and we add an edge between any two of them that are not adjacent already. This results in a graph G' of diameter p and a k -precoloring $c'_{W'}$, defined on some subset $W' \subseteq V(G')$, such that $c'_{W'}$ can be extended to a k -coloring of G' if and only if c_W can be extended to a k -precoloring of G . Moreover, the set W' forms a clique of size at most k in G' meaning that we may just as well uncolor these vertices, that is, $c'_{W'}$ can be extended to a k -coloring of G' if and only if G' is k -colorable. Hence we only need to consider:

Open Problem 6 *Determine the complexity of the problems 3-COLORING and LIST 3-COLORING for graphs of diameter 2.*

Graphs of Bounded Asteroidal Number. An *asteroidal triple* in a graph is a set of three mutually non-adjacent vertices such that each two of them are joined by a path that avoids the neighborhood of the third. An *asteroidal set* in a graph G is an independent set $S \subseteq V(G)$, such that every set of three vertices of S forms an asteroidal triple. The *asteroidal number* is the size of a largest asteroidal set in G . Note that graphs with asteroidal number at most 2 have no asteroidal triple. These graphs are also known as *AT-free* graphs. Stacho [63] proved that 3-COLORING is polynomial-time solvable on AT-free graphs. Later, Kratsch and Müller [53] extended this result by showing that even LIST k -COLORING is polynomial-time solvable on these graphs for every fixed integer $k \geq 1$. Marx [56] proved that PRECOLORING EXTENSION is NP-complete for proper interval graphs, which form a subclass of AT-free graphs. It follows from a result of Jansen [42] (see [28]) that ℓ -LIST COLORING ($\ell \geq 3$) is NP-complete for $3P_1$ -free graphs, and thus for AT-free graphs. However, the following problem, posed by Broersma et al. [7] in 1999, is still open.

Open Problem 7 *Determine the complexity of COLORING for AT-free graphs.*

Král' et al. [49] proved that COLORING is NP-complete for $4P_1$ -free graphs and thus for graphs with asteroidal number at most 3.

Open Problem 8 *Determine, for every $k \geq 3$ and $p \geq 3$, the complexity of k -COLORING, k -PRECOLORING EXTENSION and LIST k -COLORING on graphs with asteroidal number at most p .*

4 Parameterized Coloring Problems

A problem is called *fixed-parameter tractable* (FPT) if every instance (I, p) of it can be solved in time $f(p)|I|^{O(1)}$ where f is a computable function that only depends on p . If k -COLORING is polynomial-time solvable for some graph class \mathcal{G} for every integer k (and COLORING is not known to be polynomial-time solvable for \mathcal{G}) then k is a natural parameter to consider. We refer to [27] for a survey on parameterized complexity results (and open problems) with this parameter for classes of graphs characterized by one or two forbidden induced subgraphs. Here, we only mention the following open problem of Kratsch and Müller [53].

Open Problem 9 *Is COLORING fixed-parameter tractable for AT-free graphs when parameterized by k ?*

Because 3-COLORING is NP-complete in general, other parameters have been considered. For instance, Marx [55] proved that PRECOLORING EXTENSION parameterized by the number of precolored vertices is W[1]-hard for interval graphs. We survey a number of other results below (see also Table 2).

Arnborg and Proskurowski [3] proved that COLORING is FPT when parameterized by the treewidth of the input graph. Fellows et al. [21] showed that PRECOLORING EXTENSION and LIST COLORING are W[1]-hard with this parameter. On the positive side, LIST COLORING is polynomial-time solvable for any graph class of bounded treewidth, as shown by Jansen and Scheffler [44].

It is known [15] that the clique-width of a graph G is at most $2^{\text{tw}(G)-1}$, where $\text{tw}(G)$ denotes the treewidth of G . Moreover, by combining results of Kobler and Rotics [48] and Oum and Seymour [62], one finds that COLORING is polynomial-time solvable for any graph class of bounded clique-width. Hence, it is natural to research whether one can improve the FPT result for COLORING from treewidth to clique-width. However, Fomin et al. [24] showed that COLORING is W[1]-hard when parameterized by the clique-width of the input graph. We also note that PRECOLORING EXTENSION is NP-complete for distance-hereditary graphs [4], which have clique-width at most 3 [29].

The *vertex cover number* of a graph G is the size of a smallest subset $U \subseteq V(G)$, such that $G - U$ is edgeless. Fiala, Golovach and Kratochvíl [23] proved that, with this parameter, PRECOLORING EXTENSION is FPT and LIST COLORING is W[1]-hard even for split graphs. It can be observed [23] that the treewidth of a graph is at most its vertex cover number. Hence, the aforementioned result of Jansen and Scheffler [44] implies that LIST COLORING is polynomial-time solvable for any graph class of bounded vertex cover number.

The *twin cover number* of a graph G is the size of a smallest subset $U \subseteq V(G)$, such that every two adjacent vertices in $G - U$ have the same closed neighborhood in G ; note that $G - U$ is a disjoint union of cliques. Ganian [25] proved that PRECOLORING EXTENSION is FPT when parameterized by the twin cover number of the input graph. As the twin cover number of a graph is at most its vertex cover number (by definition), this result strengthens the aforementioned result of Fiala, Golovach and Kratochvíl [23]. For the same reason, LIST COLORING is W[1]-hard when parameterized by the twin cover number.

The *cluster vertex deletion number* of a graph G is the size of a smallest subset $U \subseteq V(G)$, such that $G - U$ is a disjoint union of cliques. Note that the cluster vertex deletion number of a graph G is at most its twin cover number. However, in contrast to the aforementioned FPT result of Ganian [25], Doucha and Kratochvíl [18] proved that PRECOLORING EXTENSION is W[1]-hard when parameterized by the cluster vertex deletion number of the input graph. The same authors [18] also showed that COLORING is FPT with this parameter. It is easily seen that LIST COLORING is polynomial-time solvable for any graph class \mathcal{G} of bounded cluster vertex deletion number. We guess a coloring of the set U of a graph $G \in \mathcal{G}$ that respects the given list of each vertex of U . We then

remove all vertices of U from G after adjusting the lists of the vertices in $G - U$ accordingly. As $G - U$ is a disjoint union of cliques we can solve LIST COLORING in polynomial time (see e.g. [9]). Since $|U|$ is bounded, the maximum number of colorings of U that we need to guess is polynomial. Hence, the result follows.

For an integer $c \geq 1$, the c -bounded cluster vertex deletion number of a graph G is the size of a smallest subset $U \subseteq V(G)$, such that $G - U$ is a disjoint union of cliques of size at most c . Doucha and Kratochvíl [18] proved that, for every fixed integer $c \geq 1$, PRECOLORING EXTENSION is FPT when parameterized by the c -bounded cluster vertex deletion number of the input graph.

	COLORING	PRECOLORING EXT.	LIST COL.
clique-width	W[1]-hard	para-NP-c	para-NP-c
treewidth	FPT	W[1]-hard	W[1]-hard
cluster vertex deletion number	FPT	W[1]-hard	W[1]-hard
c -bounded cluster vertex deletion number	FPT	FPT	W[1]-hard
twin cover number	FPT	FPT	W[1]-hard
vertex cover number	FPT	FPT	W[1]-hard

Table 2. The complexity of COLORING, PRECOLORING EXTENSION and LIST COLORING for various graph parameters. All problems are in XP for each parameter except when para-NP-complete. The relationships, as given in [18], between rows 1-6 are: $1 \leq 2 \leq 4 \leq 6$ and $1 \leq 3 \leq 5 \leq 6$ and $3 \leq 4$, where $x \leq y$ means that parameter x is bounded by (some function of) parameter y . Hence, membership in FPT or XP for a problem with parameter x carries over to y , and W[1]-hardness for y carries over to x .

Recently, Aboulker et al. [1] considered the number p_k of vertices of degree at least $k + 1$ of the graph G in an instance (G, k) of COLORING. This parameter is motivated by Brooks' theorem [8]: if $p_k(G) = 0$ then G is k -colorable unless G is a complete graph or an odd cycle. They showed that COLORING is FPT when parameterized by p_k .

We now discuss some graph classes that fall under the “distance from triviality” framework, introduced by Guo, Hüffner and Niedermeier [31]. For a graph class \mathcal{F} and an integer p we define four classes of “almost \mathcal{F} ” graphs, i.e. graphs that are, in some sense, “distance” p apart from \mathcal{F} , namely the classes $\mathcal{F} + pe$, $\mathcal{F} - pe$, $\mathcal{F} + pv$ and $\mathcal{F} - pv$, which consist of all graphs that can be modified into a graph of \mathcal{F} by deleting at most p edges, adding at most p edges, deleting at most p vertices and adding at most p vertices, respectively. As Grötschel, Lovász, and Schrijver, [30] proved that COLORING is polynomial-time solvable on perfect graphs, COLORING was studied from a parameterized point of view for various subclasses \mathcal{F} of perfect graphs. We survey a number of these results below. For every result mentioned, p is the chosen parameter (see also Table 3 for an overview). In this context a *modulator* of a graph is a set of at most p edges or vertices whose removal or addition makes the graph a member of \mathcal{F} .

	$+pe$	$-pe$	$+pv$	$-pv$
Bipartite	para-NP-c	P	para-NP-c	P
Chordal	FPT	FPT	W[1]-hard	P
Interval	FPT	FPT	W[1]-hard	P
Split	FPT	FPT	W[1]-hard	P
Comparability	para-NP-c	?	?	P
Complete	P	FPT	FPT	P

Table 3. The complexity of COLORING, when parameterized by p , for classes close to some subclass \mathcal{F} of perfect graphs. The polynomial and FPT cases hold even if a modulator is not part of the input.

Cai [9] proved that COLORING is FPT on split $+pe$ graphs and W[1]-hard for split $+pv$ graphs. The same author [9] also proved that whenever COLORING is polynomial-time solvable on a graph class \mathcal{F} that is closed under edge contraction and a modulator is given, then COLORING is FPT on $\mathcal{F}-pe$. As a result, COLORING is FPT for split $-pe$ graphs, and also for interval $-pe$ graphs¹ and chordal $-pe$ graphs. Note that we obtain polynomial-time solvability for $\mathcal{F}-pv$ whenever \mathcal{F} is a class of perfect graphs closed under vertex deletion (as in that case it holds that $\mathcal{F}-pv = \mathcal{F}$). Cai [9] also proved that COLORING is NP-complete for bipartite $+2v$ graphs and for bipartite $+3e$ graphs but linear-time solvable on bipartite $+1v$ and bipartite $+2e$ graphs. Marx [55] showed that COLORING is FPT² on interval $+pe$ graphs and also on chordal $+pe$ graphs but W[1]-hard for interval $+pv$ graphs and for chordal $+pv$ graphs.

An (undirected) graph is a *comparability graph* if there exists an assignment of exactly one direction to each of its edges such that (u, w) is a directed edge whenever (u, v) and (v, w) are directed edges. Takenaga and Higashide [64] proved that COLORING, restricted to comparability $+pe$ graphs, is polynomial-time solvable for $p = 1$ and NP-complete for $p \geq 2$. They also proved that COLORING is polynomial-time solvable on comparability $-1e$ graphs.

Open Problem 10 *Determine the complexity of COLORING for the classes of comparability $-pe$ graphs and comparability $+pv$ graphs when parameterized by p .*

We now consider the class of complete graphs. It is known that LIST COLORING is FPT for complete $-pe$ graphs [28]. We observe that LIST COLORING is polynomial-time solvable on complete $+pe$ graphs and complete $-pv$ graphs (because such graphs are complete and LIST COLORING is polynomial-time solvable even on block graphs [4]). In contrast to the aforementioned W[1]-hardness results of COLORING for split $+pv$, interval $+pv$ and chordal $+pv$ graphs, it holds that COLORING is FPT for complete $+pv$ graphs, as shown by Cai [9]. Golovach et al. [28] posed the following open problem.

¹ Villager et al. [66] proved afterward that a modulator can be computed in FPT time.

² For the two FPT results it was proven later, namely by Cao [10] and Marx [54], respectively, that a modulator does not have to be part of the input (but can be computed in FPT time as well).

Open Problem 11 *Determine the complexity of LIST COLORING and PRE-COLORING EXTENSION for complete+pv graphs when parameterized by p .*

Jansen and Kratsch [43] and de Weijer [69] considered the k -COLORING problem for various graph classes $\mathcal{F} + pv$ in order to obtain polynomial kernels (also some negative results are shown, for instance, 3-COLORING on path+kv has no polynomial kernel unless $\text{NP} \subseteq \text{coNP}/\text{poly}$ [43]).

5 On-line Coloring

In this section we focus on the on-line setting of graph coloring. On-line coloring algorithms were introduced by Gyárfás and Lehel [35] to model a rectangle packing problem related to dynamical storage allocation. In this setting the graph is presented vertex by vertex according to some externally determined ordering. An *on-line coloring* algorithm irrevocably colors the vertices when they come in by using a strategy that depends only on the subgraph induced by the revealed vertices and their colors. A well-known example of an on-line coloring algorithm is **First-Fit** which assigns, starting from the empty graph, each new vertex the least color from $\{1, 2, \dots\}$ that does not appear in its neighborhood. We refer to the survey of Kierstead [45] for more details.

Non-surprisingly, the number of colors used by an on-line coloring algorithm for an arbitrary graph G can be much larger than the chromatic number of G . Below we define three measures for the performance of an on-line algorithm on graphs of some specified class \mathcal{G} after first giving some additional terminology.

Let $AOL(G)$ be the (finite) set of all on-line coloring algorithms for a graph G . Let $\Pi(G)$ be the set of all permutations of the vertices of G . For $A \in AOL(G)$ and $\pi \in \Pi(G)$, let $\chi_A(G, \pi)$ denote the number of colors used by A when the vertices of G are presented to A according to π . The *A-chromatic number* $\chi_A(G)$ of G is the largest number of colors used by A to color G , that is,

$$\chi_A(G) = \max_{\pi \in \Pi(G)} \chi_A(G, \pi).$$

An algorithm A is an on-line coloring algorithm for some graph class \mathcal{G} if $A \in AOL(G)$ for every $G \in \mathcal{G}$. We let $AOL(\mathcal{G})$ be the set of on-line coloring algorithms for \mathcal{G} .

A natural performance measure for an on-line coloring algorithm, introduced by Gyárfás and Lehel [35], is to determine whether the number of colors it uses on any graph $G \in \mathcal{G}$ is bounded from above by a function that only depends on the chromatic number of G . Formally, for a graph class \mathcal{G} , we say that an algorithm $A \in AOL(\mathcal{G})$ is *competitive* if there exists a χ -*bounding* function f , that is, a function f such that

$$\chi_A(G) \leq f(\chi(G)) \text{ for every } G \in \mathcal{G}.$$

In that case \mathcal{G} is said to be *on-line χ -bounded*. For example, the class of P_4 -free graphs is on-line χ -bounded, because **First-Fit** colors every P_4 -free graph G

with $\chi(G)$ colors [35]. It is also known that **First-Fit** is competitive for interval graphs with a linear χ -bounding function [46]. Consequently, the class of interval graphs is on-line χ -bounded as well. Every class of graphs with bounded independence number [11] is also on-line χ -bounded, just as the class of P_5 -free graphs [33]. In fact, Gyárfás and Lehel [33] proved a stronger statement, namely that the class of P_5 -free graphs is *on-line ω -bounded*, that is, there exists an on-line coloring algorithm A and a function g , called an ω -bounding function, such that $\chi_A(G) \leq g(\omega(G))$ for every P_5 -free graph G . This result has been extended by Kierstead, Penrice and Trotter [47] who proved that, for every tree T of radius at most 2, the class of T -free graphs is on-line ω -bounded (with a superexponential ω -bounding function). As a special case of their result we find that the class of cocomparability graphs is on-line ω -bounded. More recently, Felsner, Micek and Ueckerdt [22] gave a cubic ω -bounding function for a subclass of cocomparability graphs, namely for the class of intersection graphs of convex sets spanned between two lines (their algorithm uses the intersection representation as input).

Despite all the above results there exist many graph classes, such as the class of trees [35], for which no competitive on-line coloring algorithm exists. These negative results lead to a natural definition of a weaker form of competitiveness, namely on-line competitiveness, which is defined as follows. The *on-line chromatic number* $\chi_{OL}(G)$ of G is the smallest number of colors used by any on-line coloring algorithm for G , that is,

$$\chi_{OL}(G) = \min_{A \in AOL(G)} \chi_A(G).$$

Then, for a graph class \mathcal{G} , an algorithm $A \in AOL(\mathcal{G})$ is said to be *on-line competitive* if there exists a function h such that

$$\chi_A(G) \leq h(\chi_{OL}(G)) \text{ for every } G \in \mathcal{G}.$$

In that case \mathcal{G} is said to be *on-line χ_{OL} -bounded*. This performance measure was coined by Gyárfás, Király, and Lehel [32], who proved that the class of graphs with girth at least 5 is on-line χ_{OL} -bounded, but results of this type have been obtained before the term was formally introduced. For instance, Gyárfás and Lehel [34] proved that, for any tree T , **First-Fit** uses $\chi_{OL}(T)$ colors.

By combining known and new results, Broersma, Capponi and Paulusma [6] proved that, for all bipartite graphs H on at most five vertices, the class of H -free bipartite graphs is on-line χ_{OL} -bounded. If H has six or more vertices the situation is not clear. For instance it is not known whether the class of C_6 -free bipartite graphs is on-line χ_{OL} -bounded. In fact this is not even known for its subclass of chordal bipartite graphs.

Open Problem 12 *Is the class of chordal bipartite graphs on-line χ_{OL} -bounded?*

Now consider P_t -free bipartite graphs. Broersma et al. [6] proved that the class of P_7 -free bipartite graphs is on-line χ_{OL} -bounded. The algorithm behind their result is based on a certain way of coloring the vertices of a complete bipartite graph with classes $\{u_1, \dots, u_m\}$ and $\{v_1, \dots, v_m\}$ minus a perfect matching $\{u_1v_1, u_2v_2, \dots, u_mv_m\}$. If the ordering is $u_1, v_1, u_2, v_2, \dots, u_m, v_m$,

then **First-Fit** assigns colors $1, 1, 2, 2, \dots, m, m$, so m colors in total. However, assigning colors $1, 1, 2, 3$ to the first four vertices u_1, v_1, u_2, v_2 in this ordering requires only three colors in total. The algorithm for P_7 -free bipartite graphs expands on this approach and uses two disjoint lists of colors for the bipartition classes of each connected component in the subgraph revealed so far. Then, whenever two connected components are glued together by an incoming vertex, it tries to prevent the “mixing” of these color lists as much as possible. Recently, Micek and Wiechert refined and extended this approach. In this way they could prove that the classes of P_8 -free bipartite graphs [58] and even P_9 -free bipartite graphs [59] are on-line χ_{OL} -bounded. This leads to the following open problem.

Open Problem 13 *Is the class of P_k -free bipartite graphs on-line χ_{OL} -bounded for every $k \geq 1$?*

As P_5 -free graphs and P_6 -free bipartite graphs are on-line ω -bounded and on-line χ_{OL} -bounded, respectively, the following problem from [6] is of interest as well.

Open Problem 14 *Is the class of (C_3, P_6) -free graphs on-line χ_{OL} -bounded?*

Solving Open Problems 12–14 may lead to new on-line χ_{OL} -bounded classes of graphs. Unlike the competitive variant, no negative results are known.

Open Problem 15 *Is the class of all graphs on-line χ_{OL} -bounded?*

6 Conclusions

We surveyed a number of results and open problems for COLORING for restricted graph classes. We did so both in an off-line and on-line setting and also considered the more general variants PRECOLORING EXTENSION and LIST COLORING. Another way of generalizing the concept of graph coloring is to consider graph homomorphisms. A *graph homomorphism* from a graph G to a graph H is a mapping $f : V(G) \rightarrow V(H)$ such that $f(u)f(v) \in E_H$ whenever $uv \in E_G$. For a fixed graph H , the problem H -HOMOMORPHISM tests whether a given graph G allows a homomorphism to H . If we choose H to be the complete graph on k vertices, then this problem is equivalent to k -COLORING. We refer to the survey of Hell and Nešetřil [37] for more on graph homomorphisms.

The classical result in the area of graph homomorphisms is the Hell-Nešetřil dichotomy theorem [38] which states that H -HOMOMORPHISM is solvable in polynomial time if H is bipartite, and NP-complete otherwise. It is a natural question whether tractability results can be obtained for non-bipartite graphs H when the input is restricted to some graph class. Not so many results are known in this direction, but to give an example, Enright, Stewart and Tardos [20] proved that, for every fixed graph H , the list version of H -HOMOMORPHISM is polynomial-time solvable for a superclass of graphs that contains the classes of permutation graphs and interval graphs.

References

1. P. Aboulker, N. Brettell, F. Havet, D. Marx and N. Trotignon, Colouring graphs with constraints on connectivity, Manuscript, arXiv:1505.01616.
2. K. Appel and W. Haken, Every planar map is four colorable, Contemporary Mathematics 89, AMS Bookstore, 1989.
3. S. Arnborg and A. Proskurowski, Linear time algorithms for NP-hard problems restricted to partial k-trees, Discrete Applied Mathematics 23 (1989) 11–24
4. F. Bonomo, G. Durán and J. Marenco, Exploring the complexity boundary between coloring and list-coloring, Annals of Operations Research 169 (2009) 3–16.
5. A. Brandstädt, T. Klemmt and S. Mahfud, P_6 - and triangle-free graphs revisited: structure and bounded clique-width, Discrete Mathematics & Theoretical Computer Science 8 (2006) 173–188.
6. H.J. Broersma, A. Capponi and D. Paulusma, A new algorithm for on-line coloring bipartite graphs, SIAM Journal on Discrete Mathematics 22 (2008) 72–91.
7. H.J. Broersma, T. Kloks, D. Kratsch and H. Müller, Independent sets in asteroidal triple-free graphs, SIAM Journal on Discrete Mathematics 12 (1999) 276–287.
8. R. L. Brooks. On colouring the nodes of a network. Mathematical Proceedings of the Cambridge Philosophical Society 37 (1941) 194–197.
9. L. Cai, Parameterized complexity of vertex coloring, Discrete Applied Mathematics 127 (2003) 415–429.
10. Y. Cao, Linear recognition of almost (unit) interval graphs, Manuscript, arXiv:1403.1515.
11. I. Cieřlik, M. Kozik, and P. Micek, On-line coloring of I_s -free graphs and co-planar graphs, Discrete Mathematics and Theoretical Computer Science Proc. AF (2006) 61–68.
12. M. Chleřík and J. Chleříková, Hard coloring problems in low degree planar bipartite graphs, Discrete Applied Mathematics 154 (2006) 1960–1965.
13. M. Chudnovsky, Coloring graphs with forbidden induced subgraphs, Proc. ICM 2014 vol IV, 291–302.
14. M. Chudnovsky, N. Robertson, P.D. Seymour, and R. Thomas, The strong perfect graph theorem, Annals of Mathematics 164 (2006) 51–229.
15. D.G. Corneil and U. Rotics, On the relationship between clique-width and treewidth, SIAM Journal on Computing 34 (2005) 825–847.
16. K. K. Dabrowski, F. Dross, M. Johnson and D. Paulusma, Filling the complexity gaps for colouring planar and bounded degree graphs, Manuscript, arXiv:1506.06564.
17. M. Demange and D. de Werra, On some coloring problems in grids, Theoretical Computer Science 472 (2013) 9–27.
18. M. Doucha and J. Kratochvřl, Cluster vertex deletion: a parameterization between vertex cover and clique-width, Proc. MFCS 2012, LNCS 7464, 348–359 (2012).
19. T. Emden-Weinert, S. Hougardy and B. Kreuter, Uniquely colourable graphs and the hardness of colouring graphs of large girth, Combinatorics, Probability & Computing 7 (1998) 375–386.
20. J. Enright, L. Stewart and G. Tardos, On list coloring and list homomorphism of permutation and interval graphs, SIAM Journal on Discrete Mathematics 28 (2014) 1675–1685.
21. M.R. Fellows, F.V. Fomin, D. Lokshtanov, F. Rosamond, S. Saurabh, S. Szeider and C. Thomassen, On the complexity of some colorful problems parameterized by treewidth, Information and Computation 209 (2011) 143–153.

22. S. Felsner, P. Micek and T. Ueckerdt, On-line coloring between two lines, Proc. SoCG 2015, LIPIcs 34 (2015) 630–641.
23. J. Fiala, P.A. Golovach and J. Kratochvíl, Parameterized complexity of coloring problems: treewidth versus vertex cover, Theoretical Computer Science 412 (2011) 2514–2523.
24. F.V. Fomin, P.A. Golovach, D. Lokshtanov and S. Saurabh, Clique-width: on the price of generality, Proc SODA 2009, 825–834 (2009).
25. R. Ganian, Twin-cover: beyond vertex cover in parameterized algorithmics. Proc. IPEC 2011, LNCS 7112 (2011) 259–271.
26. M.R. Garey, D.S. Johnson, and L.J. Stockmeyer, Some simplified NP-complete graph problems, Proc. STOC 1974, 47–63.
27. P.A. Golovach, M. Johnson, D. Paulusma and J Song, A survey on the computational complexity of coloring graphs with forbidden subgraphs, Manuscript, arXiv:1407.1482.
28. P.A. Golovach, D. Paulusma and J. Song, Closing complexity gaps for coloring problems on H-free graphs, Information and Computation 237 (2014) 204–21.
29. M.C. Golumbic and U. Rotics, On the clique-width of some perfect graph classes, International Journal of Foundations of Computer Science 11 (2000) 423–443.
30. M. Grötschel, L. Lovász, and A. Schrijver, Polynomial algorithms for perfect graphs. Annals of Discrete Mathematics 21 (1984) 325–356.
31. J. Guo, F. Hüffner and R. Niedermeier, A structural view on parameterizing problems: distance from triviality, Proc. IWPEC 2004, LNCS 3162 (2004) 162–173.
32. A. Gyárfás, Z. Király, and J. Lehel, On-line competitive coloring algorithms, Technical report TR-9703-1, 1997.
33. A. Gyárfás and J. Lehel, Effective on-line coloring of P_5 -free graphs, Combinatorica 11 (1991) 181–184.
34. A. Gyárfás and J. Lehel, First fit and on-line chromatic number of families of graphs. Ars Combinatorica 29C (1990) 168–176.
35. A. Gyárfás and J. Lehel, On-line and first-fit colorings of graphs, Journal of Graph Theory 12 (1988) 217–227.
36. P.A. Golovach and D. Paulusma, List coloring in the absence of two subgraphs, Discrete Applied Mathematics 166 (2014) 123–130.
37. P. Hell and J. Nešetřil, Graphs and Homomorphisms, Oxford Univ. Press, 2004.
38. P. Hell and J. Nešetřil, On the complexity of H -coloring, Journal of Combinatorial Theory, Series B 48 (1990) 92–110.
39. S. Huang, M. Johnson and D. Paulusma, Narrowing the complexity gap for coloring (C_s, P_t) -Free Graphs, The Computer Journal, to appear.
40. M. Hujter and Zs. Tuza, Precoloring extension. II. Graph classes related to bipartite graphs, Acta Math. Univ. Comenianae Vol. LXII (1993) 1–11.
41. M. Hujter and Z. Tuza, Precoloring extension. III. Classes of perfect graphs. Combinatorics, Probability and Computing 5 (1996) 35–56.
42. K. Jansen, Complexity Results for the Optimum Cost Chromatic Partition Problem, Universität Trier, Mathematik/Informatik, Forschungsbericht 96–41, 1996.
43. B. Jansen and S. Kratsch, Data reduction for graph coloring problems, Proc. FCT 2011, LNCS 6914 (2011) 90–101.
44. K Jansen and P. Scheffler, Generalized coloring for tree-like graphs, Discrete Applied Mathematics 75 (1997) 135–155.
45. H.A. Kierstead, Coloring graphs on-line. In: Online Algorithms: The State of the Art, LNCS 1442 (1998) 281–305.
46. H.A. Kierstead, The linearity of first-fit coloring of interval graphs, SIAM Journal on Discrete Mathematics 1 (1988) 526–530.

47. H. A. Kierstead, S. G. Penrice, and W. T. Trotter, On-line coloring and recursive graph theory. *SIAM J. Discrete Math.* 7 (1994) 72–89.
48. D. Kobler and U. Rotics, Edge dominating set and colorings on graphs with fixed clique-width, *Discrete Applied Mathematics* 126 (2003) 197–221.
49. D. Král', J. Kratochvíl, Zs. Tuza, and G.J. Woeginger, Complexity of coloring graphs without forbidden induced subgraphs, *Proc. WG 2001, LNCS 2204* (2001) 254–262.
50. J. Kratochvíl, Precoloring extension with fixed color bound. *Acta Mathematica Universitatis Comenianae* 62 (1993) 139–153.
51. J. Kratochvíl and Z. Tuza, Algorithmic complexity of list colorings, *Discrete Applied Mathematics* 50 (1994) 297–302.
52. L. Lovász, Coverings and coloring of hypergraphs, *Proc. 4th Southeastern Conf. on Combinatorics, Graph Theory, and Computing, Utilitas Math.* (1973) 3–12.
53. D. Kratsch and H. Müller, coloring AT-free graphs, *Proc. ESA 2012, LNCS 7501* (2012) 707–718.
54. D. Marx, Chordal deletion is fixed-parameter tractable, *Algorithmica* 57 (2010) 747–768.
55. D. Marx, Parameterized coloring problems on chordal graphs, *Theoretical Computer Science* 351 (2006) 407–424.
56. D. Marx, Precoloring extension on unit interval graphs, *Discrete Applied Mathematics* 154 (2006) 995–1002.
57. G.B. Mertzios and P.G. Spirakis, Algorithms and almost tight results for 3-colorability of small diameter graphs, *Algorithmica*, to appear.
58. P. Micek and V. Wiechert, An on-line competitive algorithm for coloring P_8 -free bipartite graphs, *Proc. ISAAC 2014, LNCS 8889* (2014) 516–527.
59. P. Micek and V. Wiechert, An on-line competitive algorithm for coloring bipartite graphs without long induced paths, *Manuscript*, arXiv:1502.00859.
60. M. Molloy and B. Reed, Colouring graphs when the number of colours is almost the maximum degree, *Journal of Combinatorial Theory, Series B* 109 (2014) 134–195.
61. B. Randerath and I. Schiermeyer, Vertex coloring and forbidden subgraphs - a survey, *Graphs and Combinatorics* 20 (2004) 1–40.
62. S.-l. Oum and P.D. Seymour, Approximating clique-width and branch-width, *Journal of Combinatorial Theory, Series B* 96 (2006) 514–528.
63. J. Stacho. 3-coloring AT-free graphs in polynomial time, *Algorithmica* 64 (2012) 384–399.
64. Y. Takenaga and K. Higashide, Vertex coloring of comparability+ ke and $-ke$ graphs, *Proc. WG 2006, LNCS* (2006) 102–112.
65. B. Toft and J.R. Jensen, *Graph Coloring Problems*, J. Wiley and Sons, NY, 1995.
66. Y. Villanger, P. Heggernes, C. Paul, and J.A. Telle, Interval completion is fixed parameter tractable, *SIAM Journal on Computing* 38 (2009) 2007–2020.
67. V.G. Vizing, Coloring the vertices of a graph in prescribed colors, in *Diskret. Analiz.*, no. 29, *Metody Diskret. Anal. v. Teorii Kodov i Shem* 101 (1976) 3–10.
68. K. Vušković, Even-hole-free graphs: a survey, *Applicable Analysis and Discrete Mathematics* 4 (2010) 219–240.
69. P. de Weijer, Kernelization Upper Bounds for Parameterized Graph Coloring Problems, MSc Thesis, Utrecht University, 2013.
70. Zs. Tuza, Graph colorings with local restrictions - a survey, *Discussiones Mathematicae Graph Theory* 17 (1997) 161–228.